CONTENTS OF HEREDITAS, VOLUME 140 (2004)

42 Ananias, F., Garcia, P. C. A. and Recco-Pimentel, S.
Conserved karyotypes in the *Hyla pulchella* species group (Anura, Hylidae)

134 Angus, R. B., Kemeny, C. K. and Wood, E. L.
The C-banded karyotypes of the four British species of *Notonecta* L. (Heteroptera: Notonectidae)

185 Aradottir, G. I. and Angus, R. B.
A chromosomal analysis of some water beetle species recently transferred from *Agabus* Leach to *Ilybius* Erichson, with particular reference to the variation in chromosome number shown by *I. montanus* Stephens (Coleoptera: Dytiscidae)

139 Birmeta, G., Nybom, H. and Bekele, E.
Distinction between wild and cultivated enset (*Ensete ventricosum*) gene pools in Ethiopia using RAPD markers

201 Borin, L. A. and Martins-Santos, I. C.
Study on karyotype and occurrence of B chromosomes in two endemic species of the genus *Pimelodus* (Siluriformes, Pimelodidae) from the river Iguaçu

245 Bornet, B. and Branchard, M.
Use of ISSR fingerprints to detect microsatellites and genetic diversity in several related *Brassica* taxa and *Arabidopsis thaliana*

193 Cammareri, M., Errico, A., Sebastiano, A. and Conicella, C.
Genetic relationships among *Aster* species by multivariate analysis and AFLP markers

105 Cattani, M. V. and Papeschi, A. G.
Nucleolus organizing regions and semi-persistent nucleolus during meiosis in *Spartocera fusca* (Thunberg) (Coreidae, Heteroptera)

154 Cymbron, T., Chaves, R., Adega, F., Guedes-Pinto, H. and Malheiro, M. I.
Standardization of *Msp*I and *Hae*III restriction karyotypes in cattle

149 Del Lama, M. A., Souza, R. O., Durán, X. A. A. and Soares, A. E. E.
Clinal variation and selection on MDH allozymes in honeybees in Chile

18 El Jundi, T. A. R. J. and Freitas, T. R. O.
Genetic and demographic structure in a population of *Ctenomys lami* (Rodentia-Ctenomyidae)

92 Frey, J. E. and Frey, B.
Origin of intra-individual variation in PCR-amplified mitochondrial cytochrome oxidase I of *Thrips tabaci* (Thysanoptera: Thripidae): mitochondrial heteroplasmy or nuclear integration?

229 Godfree, R. C., Woods, M. J., Young, A. G., Burdon, J. J. and Higgins, T. J.
Growth, fecundity and competitive ability of transgenic *Trifolium subterraneum* subsp. *subterraneum* cv. Leura expressing a sunflower seed albumin gene

158 Gornung, E., Mannarelli, M. E., Rossi, A. R. and Sola, L.
Chromosomal evolution in Mugilidae (Pisces, Mugiliformes): FISH mapping of the $(TTAGGG)_n$ telomeric repeat in the six Mediterranean mullets

99 Grozeva, S., Kuznetsova, V. G. and Nokkala, S.
Patterns of chromosome banding in four nabid species (Heteroptera, Cimicomorpha, Nabidae) with high chromosome number karyotypes

177 Isakova, G. K. and Mead, R. A.
Occurrence of amitotic division of trophoblast cell nuclei in blastocysts of the western spotted skunk (*Spilogale putorius latifrons*)

34 Koperdáková, J., Brutovská, R. and Čellárová, E.
Reproduction pathway analysis of several *Hypericum perforatum* L. somaclonal families

210 Ma, R., Yli-Mattila, T. and Pulli, S.
Phylogenetic relationships among genotypes of worldwide collection of spring and winter ryes (*Secale cereale* L.) determined by RAPD-PCR markers

1 Marasek, A., Hasterok, R., Wiejacha, K. and Orlikowska, T.
Determination by GISH and FISH of hybrid status in *Lilium*

70 Nkongolo, K. K., Kim, N. S. and Michael, P.
Detection and physical mapping of the 18S-5.8S-26S rDNA and the pKFJ660 probe with microsatellite sequences derived from the rice blast fungus (*Magnaporthe grisea*) in conifer species

123 Offei, S. K., Asante, I. K. and Danquah, E. Y.
Genetic structure of seventy cocoyam (*Xanthosoma sagittifolium*, Linn, Schott) accessions in Ghana based on RAPD

129 O'Leary, N., Calviño, C. I., Greizerstein, E., Martínez, S. and Poggio, L.
Further cytogenetical studies on diploid and polyploid species of *Eryngium* L. (Saniculoideae, Apiaceae) from Argentina

24 Oliveira, K. M., Laborda, P. R., Garcia, A. A. F., Zagatto Paterniani, M. E. A. G. and Souza, A. P.
Evaluating genetic relationships between tropical maize inbred lines by means of AFLP profiling

81 Ortiz, R. and Vorsa, N.
Transmission of a cyclical translocation in two cranberry cultivars

87 Rodrigues, L. R. R., Barros, R. M. S., Pissinati, A., Pieczarka, J. C. and Nagamachi, C. Y.
A new karyotype of an endangered primate species (*Callicebus personatus*) from the Brazilian Atlantic forests

161 Rožek, M., Lachowska, D., Petitpierre, E. and Holecová, M.
C-bands on chromosomes of 32 beetle species (Coleoptera: Elateridae, Cantharidae, Oedemeridae, Cerambycidae, Anthicidae, Chrysomelidae, Attelabidae and Curculionidae)

250 Sazanov, A. A., Sazanova, A. L., Tzareva, V. A., Kozyreva, A. A., Smirnov, A. F., Romanov, M. N., Price, J. A. and Dodgson, J. B.
Chromosomal localization of three GGA4 genes using BAC-based FISH mapping: a region of conserved synteny between the chicken and human genomes

61 Sharma, P. N., Torii, A., Takumi, S., Mori, N. and Nakamura, C.
Marker-assisted pyramiding of brown planthopper (*Nilaparvata lugens* Stål) resistance genes *Bph*1 and *Bph*2 on rice chromosome 12

49 Šťáhlavský, F. and Král, J.
Karyotype analysis and achiasmatic meiosis in pseudoscorpions of the family Chthoniidae (Arachnida: Pseudoscorpiones)

171 Tiidema, A. and Truve, E.
Efficient regeneration of fertile barley plants from callus cultures of several Nordic cultivars

222 Toker, C.
Estimates of broad-sense heritability for seed yield and yield criteria in faba bean (*Vicia faba* L.)

226 Toker, C. and Ilhan Cagirgan, M.
The use of phenotypic correlations and factor analysis in determining characters for grain yield selection in chickpea (*Cicer arietinum* L.)

112 Veličković, M.
Chromosomal aberrancy and the level of fluctuating asymmetry in black-striped mouse (*Apodemus agrarius*): effects of disturbed environment

8 Wang, T., Su, Y.-J., Li, X.-Y., Zheng, B., Chen, G.-P. and Zeng, Q.-L.
Genetic structure and variation in the relict populations of *Alsophila spinulosa* from southern China based on RAPD markers and cpDNA *atp*B-*rbc*L sequence data